

Influence of the left-handed part of the neutrino mass matrix on the lepton number violating $e^-e^- \rightarrow W^-W^-$ process

P. Duka, J. Gluza, M. Zralek

*Department of Field Theory and Particle Physics Institute of Physics, University of Silesia Uniwersytecka 4, PL-40-007
Katowice, Poland E-mails: duka,gluza,zralek@us.edu.pl*

(February 1, 2008)

Influence of the neutrino mass submatrix M_L on the $e^-e^- \rightarrow W^-W^-$ process is discussed. Taking into account various possible CP signatures of heavy neutrinos it is shown that, in some cases, nonzero M_L substantially changes predictions for maximum possible values of the $e^-e^- \rightarrow W^-W^-$ cross section. A direct role of the ω^2 parameter (coming from neutrinoless double beta decay) is clarified. The consequences of doubly charged Higgs particles (δ^{--}) with resonances even far away from energies of the future linear lepton collider ($\sqrt{s} = 0.5 \div 1$ TeV) are studied.

13.15.-f, 12.15.Cc, 11.30.Er

I. INTRODUCTION

The e^-e^- option of the future $\sqrt{s} = 0.5 \div 2$ TeV linear collider is a very interesting area for investigating new physics [1]. Processes like $e^-e^- \rightarrow l_i^- l_j^-$, W^-W^- ($l_{i(j)} = e, \mu, \tau$) could be studied indicating that lepton number (flavour or total) is not a global symmetry of the electroweak interactions. In this paper we would like to examine the $e^-e^- \rightarrow W^-W^-$ process. This reaction violates the total lepton number by two units, $\Delta L = 2$. Its potential importance and hopes connected with it are based on two facts. First, the SM background is very small and under control [2] so with the planned luminosity $10 \text{ fb}^{-1}/\text{year}$ [1] the cross section as small as 0.1 fb could give visible effect. Second, its occurrence would indicate that there exist massive neutrinos of Majorana type. These neutrinos must be heavy (with masses $M_N > M_Z$) as known neutrinos cannot give any substantial signal [3]. Many papers have been devoted to this process during the last decade [4–7]. For the first time this reaction was proposed and examined in 1982 by Rizzo [4]. Additional interest has come with the paper [5] where it was shown that the process is enhanced for heavy neutrino masses in the vicinity of the collider's c.m. energy. Then the optimism was revised in [6] where constraints on heavy neutrinos coming from neutrinoless double- β decay have been taken into account. It has been shown that an observable signal requires fine-tuning among different heavy neutrino couplings. However, as it has been shown in [8], these cancelations can be in a natural manner connected with CP parities of heavy neutrinos. All other papers cited in [7] give many interesting details connected with the process.

This paper brings another such a detail which can however appear to be crucial for the magnitude of the cross section. In the last paper concerning the $e^-e^- \rightarrow W^-W^-$ where all relevant constraints on the heavy neutrinos have been taken into account [8] we have assumed that the neutrino mass submatrix M_L generated by left-handed neutrinos

$$M = \begin{pmatrix} M_L & M_D \\ M_D^T & M_R \end{pmatrix}, \quad (1)$$

is exactly zero (M_D is the submatrix of Dirac type masses, M_R is the submatrix of right-handed Majorana masses). This means that we have considered the class of models beyond the standard where only right-handed neutrinos were introduced. Then, to get an observable magnitude of the cross section, at least 3 heavy neutrinos with appropriate CP signatures and masses were necessary. However, there are models where M_L does not vanish. Such nonzero M_L changes the relations which restrict the space of parameters of possible (i.e. allowed by experimental data) heavy neutrinos couplings and masses.

The full phenomenological discussion of non-zero M_L has been given lately [9] in the context of heavy neutrino production in e^-e^+ ($e^-e^+ \rightarrow \nu N$) and $e^- \gamma$ ($e^- \gamma \rightarrow W^- N$) reactions. Here we will restrict ourselves to two non-standard models with possible nonzero M_L . The Standard Model with both additional right handed neutrinos (RHS) and Higgs triplets and the Left-Right symmetric model (LR). Details of these models can be found in literature (e.g. in [3,10]). As we are going to find the largest possible values of the cross section we consider models where CP is conserved in the lepton sector.

II. THE INFLUENCE OF M_L ON $E^-E^- \rightarrow W^-W^-$.

The leading helicity amplitudes for the $e^-e^- \rightarrow W^-W^-$ process can be written in the following, simplified, way

$$M = \sum_a \left\{ K_{ae}^2 m_a [f_t(m_a) + f_u(m_a) + f_s^L] + (K_R)_{ae}^2 m_a f_s^R \right\}. \quad (2)$$

The matrices K, K_R are part of the unitary matrix $U = (K^*, K_R)^T$ which diagonalizes the 6×6 neutrino mass matrix M in Eq.(1), the index L(R) is connected with the left (right) doubly charged Higgs particle which is exchanged in the s channel. For details see e.g. [3], [10].

The sum in Eq.(2) runs over all light (ν) and heavy (N) neutrinos. Let's note that the kinematical factors in t and u channels $f_{t,u}$ depend on m_a , but the $f_s^{L(R)}$ ones in s channel do not. First we will examine the t and u channels assuming only that the influence of the s channel is negligible (heavy $\delta_{L,R}^-$). At the end we will comment on the effect of non-zero M_L on the s-channel contribution.

The experimental bounds on the elements of the matrix, $K_{\nu e}$ and K_{Ne} , describing the mixing of electrons with light and heavy neutrinos can be summarized as follows

$$\sum_{N(heavy)} |K_{Ne}^2| \leq \kappa^2 = 0.0054, \quad (3)$$

$$\sum_{\nu(light)} |K_{\nu e}^2 m_\nu| \leq \kappa_{light}^2 = 0.65 \text{ eV}, \quad (4)$$

$$\left| \sum_{N(heavy)} K_{Ne}^2 \frac{1}{m_N} \right| \leq \omega^2 = 6 \cdot 10^{-3} \div 5 \cdot 10^{-5} \text{TeV}^{-1}. \quad (5)$$

The first relation (Eq.(3)) comes from low energy experiments [11], the other ones can be derived from the fact that neutrinoless double- β decay $(\beta\beta)_{0\nu}$ has not been detected yet¹.

Diagonalization of the matrix (1) together with (4) yields to the following relation ($m_L = (M_L)_{\nu_e \nu_e}$)

$$\left| m_L - \sum_N K_{Ne}^2 m_N \right| < \kappa_{light}^2. \quad (6)$$

However, κ_{light}^2 is very small and can be neglected, then from (6) we get

$$\sum_N m_N K_{Ne}^2 = m_L. \quad (7)$$

Similar to the analysis given in [8,9] let us discuss the influence of m_L on the magnitude of the cross section for different CP parities of heavy neutrinos.

If we have only one heavy neutrino state (or more but with the same CP parities) then from (3),(5) and (7) we get restrictions on m_L [9]

$$0 \leq m_L \leq \min(\kappa^2 M, \omega^2 M^2), \quad (8)$$

where M is the mass of the lightest of heavy neutrinos. It gives for instance $m_L \leq 5 \cdot 10^{-4} \text{ GeV}$ for $M=100 \text{ GeV}$ ($\omega^2 = 5 \cdot 10^{-5} \text{ TeV}^{-1}$). For the above values of m_L the mixing angle K_{Ne} is limited to [9]

$$K_{Ne}^2 \leq \min(\omega^2 M, \kappa^2). \quad (9)$$

¹As we can see there exist large discrepancies in the limit on ω^2 . For arguments on lower (upper) limits, see [12] ([13]).

Fig.1 shows the maximum value of the cross section $\sigma(e^-e^- \rightarrow W^-W^-)$ where the parameters are restricted by relation (9) for three different values of ω^2 and $\sqrt{s} = 0.5(1)$ TeV. We can see that for various ω^2 there are different masses M_0 for which the cross section reaches maximum value, e.g. $M_0 \simeq 1(100)$ TeV for $\omega^2 = 6 \cdot 10^{-3}$ ($5 \cdot 10^{-5}$) TeV^{-1} . For $M \leq M_0$ the maximum value of the cross section increases with increasing M ($(K_{Ne})_{max}^2$ in Eq.(9) increases), for masses larger than M_0 the cross section decreases with M ($(K_{Ne})_{max}^2 = \kappa^2 = \text{const}$). We can see that only for $\omega^2 > 5 \cdot 10^{-4} TeV^{-1}$ and $\sqrt{s} \geq 1$ TeV there is a small region of masses where $\sigma_{max} > 0.1$ fb. If there is only one heavy neutrino or more but with the same CP parities then the value of ω^2 crucially determines σ_{max} . Much effort is devoted to find the bound on ω^2 parameters [12].

For the case of two heavy neutrinos with opposite CP parities we get the following inequalities ($K_{N_1e} = x_1$, $K_{N_2e} = ix_2$, $m_1 = M$, $m_2 = AM$)

$$x_1^2 + \left| \frac{x_1^2}{A} - \frac{m_L}{AM} \right| \leq \kappa^2, \quad (10)$$

$$\left| x_1^2 \left(1 - \frac{1}{A^2} \right) + \frac{m_L}{A^2 M} \right| \leq \omega^2 M. \quad (11)$$

When $m_L = 0$, to remove the bound given by ω^2 (Eq.(11)), we have to assume that two neutrinos are almost degenerate: $A \rightarrow 1$. But then we have practically one Dirac neutrino (two Majorana neutrinos with opposite CP values) and the cross section approaches zero. This was actually shown in [8] where the $m_L = 0$ case was examined.

However, for $m_L \neq 0$ situation is different. The inequalities (10) and (11) can be satisfied only for the confined region of m_L [9]

$$- \max(AM\kappa^2, (A-1)M\kappa^2 + A^2\omega^2 M^2) \leq m_L \leq \min(\omega^2 M^2, \kappa^2 M). \quad (12)$$

Positive values of m_L are strongly restricted but the space of negative m_L values is wider and depends on the values of M and A . In Fig.2 we plot the results for $m_L = -1(-3, -5)$ GeV and $A=5$ as a function of neutrino mass. As we can see lines start from different masses. This is because Eq.(12) must holds. Similar results can be obtained for larger spectrum of A ($= 3 \div 15$). For positive m_L the situation is similar to the case with $n_R = 1$ (compare Eqs.(8),(12)). Results given in Fig.2 describe also the case of three heavy neutrinos with the following CP signatures $\eta_{CP}(N_1) = -\eta_{CP}(N_2) = -\eta_{CP}(N_3)$. Then two heavy neutrinos (N_2, N_3) contribute in the same way to the amplitude (Eq.(2)) and can be effectively treated as one.

The last quantitatively distinguishable possibility which is left for three heavy neutrinos is the case $\eta_{CP}(N_1) = \eta_{CP}(N_2) = -\eta_{CP}(N_3)$. Then initial inequalities (3,5) and (7) are satisfied if m_L is confined to the following region ($m_1 = M$, $m_2 = AM$, $m_3 = BM$) [9]

$$\begin{aligned} - \min \{ BM\kappa^2, \max \{ B^2\omega^2 M^2, (B-1)M\kappa^2 + B\omega^2 M^2 \} \} &\leq m_L \\ &\leq \min \{ AM\kappa^2, (A-B)M\kappa^2 + AB\omega^2 M^2 \}. \end{aligned} \quad (13)$$

By fixing $B=10$ and $M=100$ GeV for different values of A we have found mixing angles $K_{N_1e} = x_1$, $K_{N_2e} = x_2$, $K_{N_3e} = ix_3$ such that the cross section is maximal. The result is given in Fig.3 for $\sqrt{s} = 1$ TeV. For larger masses ($M > 100$ GeV) σ_{max} decreases, e.g. for $M=200$ GeV, $\sigma_{max} \leq 4$ fb. Let us note that the largest results are possible for large A and then we can always find a space of allowed mixings for which $\sigma_{max} \simeq 9$ fb independently of m_L . Similar plots can be made for other energies $0.5 TeV \leq \sqrt{s} \leq 2$ TeV with a result $\sigma_{max} \leq 1(25, 40)$ fb and $\sqrt{s} = 0.5(1.5, 2)$ TeV, respectively (see [8] for the $m_L = 0$ case).

Finally, in Fig.4 we describe the s-channel contribution to the $e^-e^- \rightarrow W^-W^-$ process. We present the contribution of two doubly charged Higgs particles δ_L^{--} and δ_R^{--} which exist for example in the LR model². Masses of the $\delta_{L,R}^{--}$ particles depend on M_{W_2} [15] and for $M_{W_2}=1$ TeV we have (without fine tuning between parameters in the Higgs potential) $M_{\delta_L^{--}} \simeq 1600$ GeV and $M_{\delta_R^{--}} \simeq 3000$ GeV. As $m_{\delta_R^{--}} \gg m_{\delta_L^{--}}$ the effect of δ_R^{--} is negligible. In such circumstances our considerations are also valid for the SM enlarged by additional Higgs triplet and right handed neutrinos. Let us note that the contribution of the δ_L^{--} resonance to the helicity amplitudes (Eq.(2)) is directly proportional to m_L ($m_L = \sum_a K_{ae}^2 m_a$, see e.g. [10]) and is invisible if only light neutrinos exist. If we take $m_L = 0$ then the resonance disappears (solid line in Fig.4). If however $m_L \neq 0$ then its effect can be large. This is shown in

²Other aspects of doubly charged Higgs physics at an e^-e^- collider can be found in [14]

Fig.4 where we take doubly charged Higgs' widths to be $\Gamma_{\delta_{L,R}^{--}} = \Gamma_{M_W} M_{\delta_{L,R}^{--}} / M_W$. Lines on this Figure present the cross sections for the case when all heavy neutrinos have the same CP eigenvalues. As it has already been discussed, σ_{max} depends strongly on ω^2 in this case. We take $\omega^2 = 5 \cdot 10^{-5} \text{ TeV}^{-1}$, so t and u channels contributions to the cross section are very small (see Fig.1). This means that the large cross sections in Fig.4, even for energies far away from resonance region, are due to the δ_L^{--} resonance. For example for $m_L = 5 \text{ GeV}$ and $\sqrt{s} = 1 \text{ TeV}$ the $\sigma_{max} \simeq 40 \text{ fb}$ and the effect is caused almost exclusively by $\delta_L^{--}(1600)$ Higgs resonance. As the contribution of δ_R^{--} to the cross section is not proportional to m_L its effect can be large even for $m_L = 0$ especially if its mass is around the C.M. energy. This case has been considered in [16].

This is shown explicitly in Fig.5 where s-channel contribution to the $e^-e^- \rightarrow W^-W^-$ process as function of energy is presented. To extract the effect of the δ_L^{--} resonance we compare the cross section σ_{max} for t and u channels only (short-dashed line) with the total cross section where t,u and s channels are added altogether (long-dashed lines) for $m_L = 1 \text{ GeV}$ and $\eta_{CP}(N_1) = \eta_{CP}(N_2)$. We can see the huge influence of δ_L^{--} resonance on the total cross section. Even for very high mass of δ_L^{--} ($M_{\delta_L^{--}} = 2000 \text{ GeV}$, $\sqrt{s} = 1 \text{ TeV}$) σ_{max} is above the "discovery limit".

Solid lines in Fig.5 describe another case with $\eta_{CP}(N_1) = -\eta_{CP}(N_2)$. The upper one corresponds to the full cross section (s,t,u channels), the lower one is for a cross section without the s channel. As we can see for $m_L = -1 \text{ GeV}$, contributions of the s and t+u channels are now comparable. The influence of δ_L^{--} on the cross section depends on the δ_L^{--} mass and width, and the value of the m_L parameter. For the same mass and the same width of δ_L^{--} its contribution to $\sigma_{max}(e^-e^- \rightarrow W^-W^-)$ can be very small, comparable or much bigger than the t+u channels' part depending on the value of m_L .

III. CONCLUSIONS

We have analyzed the predictions for maximum possible value of the $e^-e^- \rightarrow W^-W^-$ cross section in models with nonzero m_L . If we have only one heavy neutrino or more but with the same CP parities then the value of ω^2 is crucial for the maximum of the cross section and m_L does not have any visible influence. For the smallest value of ω^2 ($\leq 5 \times 10^{-5} \text{ TeV}^{-1}$) predicted by some existing estimations (e.g. [13]) the cross section σ_{max} is too small to be measured in the future e^-e^- linear colliders unless $\delta_{L,R}^{--}$ exist in the model.

However, for all other cases, the $m_L \neq 0$ changes substantially the $e^-e^- \rightarrow W^-W^-$ cross section. Negative m_L values move the limits on experimentally allowed neutrino mixings and masses. If there are two heavy neutrinos with opposite CP parities (or any number of them but with the lightest one having opposite CP parity with respect to all other ones) the value of σ_{max} can be substantial, much above the background level (e.g. for $M=150 \text{ GeV}$, $A=5$, $\sigma_{max} \simeq 7 \text{ fb}$).

In another configuration of CP parities of heavy neutrinos ($\eta_{CP}(N_1) = \eta_{CP}(N_2) = -\eta_{CP}(N_3)$) the largest $\sigma(e^-e^- \rightarrow W^-W^-)$ is obtained for $m_L = 0$.

The most dramatic influence of the non-zero m_L on $e^-e^- \rightarrow W^-W^-$ is connected with the δ_L^{--} resonance. For $m_L = 0$ the contribution of this resonance to the process disappears. The $m_L \neq 0$ values cause the δ_L^{--} to give large contribution even far away from on-peak energies. The contribution of the δ_R^{--} to the cross section does not depend on the m_L value.

ACKNOWLEDGMENTS

This work was supported by Polish Committee for Scientific Research under Grant No. PB659/P03/95/08 and University of Silesia internal Grant.

J.G. also appreciate the fellowship of the Foundation for Polish Science.

-
- [1] Physics and technology of the Next Linear Collider: a report submitted to Snowmass '96 by NLC ZDR Design Group and NLC Physics Working Group (S. Kuhlman et al.), SLAC-R-0485, hep-ex/9605011.
 - [2] V. Barger, J.F. Beacom, K. Cheung, T. Han Phys. Rev. **D50** (1994) 6704 and **D52** (1995) 3815; J.F. Gunion and A. Tofighi-Niaki, Phys. Rev. **D36** (1987) 2671 and **D38** (1988) 1433; F. Cuyper, K. Kolodziej, O. Korakianitis and R. R  kl Phys. Lett. **B325** (1994) 243.

- [3] J. Gluza and M. Zralek, Phys. Lett. **B362**(1995)148.
- [4] T. Rizzo, Phys.Lett.**B116** (1982)23.
- [5] C.A. Heusch and P. Minkowski, Nucl. Phys. **B416** (1994) 3.
- [6] G. Belanger, F. Boudjema, D. London and H. Nadeau, Phys. Rev. **D53** (1996) 6292.
- [7] D. London, G. Belanger and J.N. Ng, Phys.Lett. **B188** (1987)155; J. Maalampi, A. Pietila and J. Vuori, Nucl.Phys. **B381** (1992)544; C.A. Heusch and P. Minkowski, Phys.Lett.**B374**(1996)116; P. Helde, K. Huitu, J. Maalampi, M. Raidal, Nucl.Phys. **B437** (1995)305; T. Rizzo Int. J. Mod. Phys. **A11**(1996)1563 and 1613; C.A. Heusch and P. Minkowski, SCIPP-96-41, hep-ph/9611353; B. Ananthanarayan and P. Minkowski, Phys. Lett. **B373**(1996)130.
- [8] J. Gluza and M. Zralek, Phys. Lett. **B372** (1996) 259.
- [9] J. Gluza, J. Maalampi, M. Raidal and M. Zralek, Phys. Lett. **B407** (1997) 45.
- [10] J. Gluza and M. Zralek, Phys. Rev. **D52** (1995) 6238.
- [11] See e.g., A. Djouadi, J. Ng and T.G. Rizzo, hep-ph/9504210; P. Langacker and D. London Phys. Rev. **D38** (1988) 886; E. Nardi, E. Roulet, D. Tommasini Nucl. Phys. **B386** (1992) 239; Phys. Lett. **B344** (1995) 225; C.P. Burgess et al. Phys. Rev. **D49** (1994) 6115.
- [12] C. Greub and P. Minkowski, hep-ph/9612340.
- [13] See e.g., J.D. Vergados, Phys. Rev. **D28** (1983) 2887; Phys. Rep. **133** (1986) 1; A. Tomada, Rep. Prog. Phys. **54** (1991) 1; T. Bernatowicz et al., Phys. Rev. Lett. **69** (1992) 2341; A. Balysh et al. (Heidelberg-Moscow Coll.), Phys. Lett. **B356** (1995) 450; M. Hirsch, H.V. Klapdor-Kleingrothaus and O. Panella, Phys. Lett. **B374**(1996)7.
- [14] J.F. Gunion, Int. J. Mod. Phys. **A11**(1996)1551; J.F. Gunion, C. Loomis, K.T. Pitts, hep/ph/9610237.
- [15] J. Gluza and M. Zralek, Phys.Rev.**D51**(1995)4695.
- [16] J. Gluza, Phys.Lett.**B403**(1997)304.




FIG. 1. The largest cross section for the $e^+e^- \rightarrow W^+W^-$ process with any number of heavy neutrinos with the same CP parities. Dashed (solid) line is for $\sqrt{s} = 0.5(1)$ TeV energy, (1),(2),(3) stand for different ω^2 values: $\omega^2 = 6 \cdot 10^{-3} \text{ TeV}^{-1}$ (1); $\omega^2 = 5 \cdot 10^{-4} \text{ TeV}^{-1}$ (2); $\omega^2 = 5 \cdot 10^{-5} \text{ TeV}^{-1}$ (3). Doubly solid line in this and next figures denotes a background level of this process.




FIG. 2. Influence of the m_L on σ_{max} for two heavy neutrinos with opposite CP parities for $\sqrt{s} = 1 \text{ TeV}$ and $A=5$ as function of M . Only negative m_L values give substantial results in this case.




FIG. 3. The case of three heavy neutrinos with $\eta_{CP}(N_1) = \eta_{CP}(N_2) = -\eta_{CP}(N_3)$. The cross section as function of m_L for different A , $B=10$ and $M=100$ GeV is given. (a),(b),(c),(d),(e) are for $A = 10^6, 100, 50, 20, 10$, respectively.




FIG. 4. Influence of the m_L parameter on the s-channel δ_L^{--} resonance. Solid, dashed and dotted lines are for $m_L = 0, 1, 5$ GeV, respectively. The t and u channel contributions are calculated for the same η_{CP} eigenvalues of heavy neutrinos and $\omega^2 = 5 \cdot 10^{-5} \text{ TeV}^{-1}$.




FIG. 5. Contribution of the δ_L^{--} resonance to the $e^-e^- \rightarrow W^-W^-$ process in the range of energies of the future linear lepton collider. Dashed lines are for $m_L = 1$ GeV and the same CP parities of heavy neutrinos (case A), solid lines are for $m_L = -1$ GeV and $\eta_{CP}(N_1) = -\eta_{CP}(N_2)$ (case B). To show the s channel effect we present σ_{max} for t and u channels only (short dashed line for the A case and lower solid line for the B case) and for the full cross section with s,t and u channels altogether (long dashed lines for the A case and upper solid line for the B case). Long dashed lines are for $M_{\delta_L^{--}} = 1000, 1600, 2000$ GeV, respectively. The upper solid line is for $M_{\delta_L^{--}} = 1600$ GeV.